arXiv:1610.08046v1 [cond-mat.stat-mech] 24 Oct 2016

One-Dimensional Random Walk in Multi-Zone Environment

AV Nazarenko[1] and V Blavatska[2]

[1] Bogolyubov Institute for Theoretical Physics of the National Academy of Sciences of Ukraine, UA–03680 Kyiv, Ukraine

[2] Institute for Condensed Matter Physics of the National Academy of Sciences of Ukraine, UA–79011 Lviv, Ukraine

E-mail: nazarenko@bitp.kiev.ua

Abstract. We study a symmetric random walk (RW) in one spatial dimension in environment, formed by several zones of finite width, where the probability of transition between two neighbouring points and corresponding diffusion coefficient are considered to be differently fixed. We derive analytically the probability to find a walker at the given position and time. The probability distribution function is found and has no Gaussian form because of properties of adsorption in the bulk of zones and reflection at the separation points. Time dependence of the mean squared displacement of a walker is studied as well and revealed the anomalous behaviour as compared with ordinary RW.

Keywords: Random Walk, Inhomogeneous Environment, Diffusion

PACS numbers: 05.40.Fb, 02.50.Ga

1. Introduction

The model of random walk (RW) finds its application in the majority of fields like polymer physics, economics, computer sciences [1] and is generally used as simple mathematical formulation of diffusion process. The standard unconstrained RW is characterized by constant transition probabilities which are independent in space and time, being the prototype example of a stochastic Gaussian process.

The RW problem in inhomogeneous environment is of great interest since it is connected with physical phenomena like transport properties in fractures and porous rocks, diffusion of particles in gels, colloidal solutions and biological cells (see, e.g., [2] for a review). Note that restricting the volume of space available to the diffusing particles by confining boundaries causes remarkable entropic effects [3].

Numerous studies, dealing with the one-dimensional RW in environment with "inhomogeneities" such as walls [4, 5, 6], stack of permeable barriers [7, 8, 9] or finite-sized barriers [10, 11], revealed the physically significant effects and the non-trivial behaviour with pronounced deviation away from Gaussian one. They also point out the difficulties in analytical description in the problems above. Taking into account the interesting properties of such models, we are aiming to generalize them.

In the present paper, we focus our attention on the symmetric RW model with position dependent transition probability which is equal for the left and right steps and varies within the different intervals (zones) of coordinate axis. Actually, replacing the probabilities of transition and adhesion within each zone with their mean values, we receive the step-like dependence in space. In particular, the zones of narrow width in our model reproduce the walls.

Thus, we consider RW in environment, consisting of N zones with constant parameters, and reduce the problem to solving the differential equation with diffusion coefficient which inherits the established space dependence. We concentrate our attention on finding the probability distribution function (PDF) by neglecting the local fluctuations of environment characteristics. Since the adhesion property of zones can vary, it is expected that a time dependence of the averages should differ from that in the case of uniform medium.

A particle performing ordinary diffusion is typically characterized by the mean square displacement $\langle x^2 \rangle$ obeying a scaling law $\langle x^2 \rangle \sim t$ at time t (or, equivalently, after t steps of RW trajectory) independently of the space dimension d [12]. Usually, presence of inhomogeneities causes the anomalous diffusion regime with non-linear power-law dependence on t. On the basis of PDF found for any N we calculate $\langle x^2 \rangle$ as function of finite t and investigate its anomalous behaviour within the triple-zone models.

Averaging PDF over time t, we can also estimate a probability to find a walker in a given point of space and compare it with the performed numerical simulations. We interpret its profile peculiarities from the points of view of transport theory and mechanics.

The layout of the paper is as follows. In the next Section, we constitute the

random walk rules and obtain the differential equation for PDF. The probability function and its distribution (PDF) are found in Section 3. After calculating the mean square displacement in Section 4, we end up with giving discussion and outlook.

2. Formulation of RW Model in N-Zone Environment

We start with discrete RW model, which can be considered as a Markov process, where unitary step in time results in either zeroth or unitary step in space. The evolution is determined here by stationary transition probability $T(x_{t+1}, x_t)$ defined as

$$T(x,y) = \begin{cases} p(x)\delta_{x-y,-1} + q(x)\delta_{x-y,1} + r(x)\delta_{x-y,0}, & \text{if } |x-y| \le 1 \\ 0, & \text{if } |x-y| > 1, \end{cases} \tag{1}$$

where constraint $p(x)+q(x)+r(x) = 1$ is fulfilled for all x; $\delta_{x,y}$ is the Kronecker symbol for $x \in \mathbb{Z}$.

Functions $p(x)$ and $q(x)$ determine probabilities to find a walker at point x if it was at $x+1$ or $x-1$ in previous time moment, respectively; while $r(x)$ corresponds to probability of adhesion (adsorption) [5]. In general, $p(x)$, $q(x)$ are regarded as arbitrary non-negative functions which do not exceed the unit.

To formulate our model, we give the finite sequence $\{a_n\}$ of positive numbers,

$$0 = a_0 < a_1 < \ldots < a_{N-1} < a_N = \infty, \tag{2}$$

determining the positions of walls or separation points of environment phases.

For $x < 0$, we reproduce the same configuration symmetric with respect to the starting point.

Using the finite set $\{p_n\}$ of the given positive constants, we consider RW without preference between the moves to the left and right in heterogeneous environment with the following properties:

$$p(x) = q(x) = p_n, \qquad r(x) = 1 - 2p_n, \quad \text{for} \quad a_{n-1} < |x| < a_n; \tag{3}$$

$$p(a_n) = q(-a_n) = p_{n+1}, \quad q(a_n) = p(-a_n) = p_n, \tag{4}$$

$$r(\pm a_n) = 1 - p_n - p_{n+1}. \tag{5}$$

To preserve a probability meaning of the functions, we require $p_n \le 1/2$.

Time evolution of the probability distribution function (PDF) $P(x,t)$ in Markov picture with our initial condition $P(x,0) = \delta_{x,0}$ is given by equation:

$$P(x,t+1) = \sum_{y\in\mathbb{Z}} T(x,y)P(y,t), \qquad \sum_{y\in\mathbb{Z}} P(y,t) = 1. \tag{6}$$

Master equation with arbitrary distance ℓ and time τ between successive steps reads

$$P(x,t+\tau) = p(x)P(x+\ell,t) + q(x)P(x-\ell,t) + r(x)P(x,t). \tag{7}$$

To obtain a differential equation for PDF, t should be relatively large while ℓ and τ tend to zero simultaneously. Using a Taylor expansion of $P(x,t)$, one gets

$$\partial_t P(x,t) = \upsilon(p(x)-q(x))\partial_x P(x,t) + \mathcal{D}(p(x)+q(x))\partial_x^2 P(x,t) + O(\ell^2), \tag{8}$$

here $\upsilon \equiv \ell/\tau$ and $\mathcal{D} \equiv \ell^2/2\tau$ are constant drift velocity and diffusivity, respectively.

Omitting terms of order $O(\ell^2)$ and fixing $\upsilon = 1$ and $\mathcal{D} = 1/2$, we obtain differential equation of RW in diffusion approximation:

$$\partial_t P(x,t) = \mathrm{v}(x)\partial_x P(x,t) + D(x)\partial_x^2 P(x,t), \tag{9}$$

where $D(x) = p_n$ and $\mathrm{v}(x) = 0$ for $a_{n-1} < |x| < a_n$; $D(\pm a_n) = (p_n + p_{n+1})/2$ and $\mathrm{v}(\pm a_n) = \pm(p_{n+1} - p_n)$ for $n = \overline{1, N-1}$.

We can formally write

$$\mathrm{v}(x) = \mathrm{sign}(x)\sum_{n=1}^{N-1}\alpha_n\delta_n(x), \qquad D(x) = \sum_{n=1}^{N} p_n\chi_n(x), \tag{10}$$

where $\alpha_n = p_{n+1} - p_n$, $|\alpha_n| < 1/2$; and

$$\delta_n(x) = \begin{cases} 1, & |x| = a_n \\ 0, & \text{otherwise} \end{cases}, \tag{11}$$

$$\chi_1(x) = \begin{cases} 1, & |x| < a_1 \\ 1/2, & |x| = a_1 \\ 0, & \text{otherwise} \end{cases}, \quad \chi_{n>1}(x) = \begin{cases} 1, & a_{n-1} < |x| < a_n \\ 1/2, & |x| = a_{n-1}, a_n \\ 0, & \text{otherwise} \end{cases}. \tag{12}$$

Characteristic functions $\{\chi_n\}$ of intervals are orthogonal and have the properties:

$$\sum_{n=1}^{N}\chi_n(x) = 1, \quad |x| < \infty; \qquad \chi_n(x)\chi_m(x) = 0, \quad n \neq m; \tag{13}$$

$$\int_{-\infty}^{\infty}\chi_n(x)\chi_m(x)\mathrm{d}x = 2(a_n - a_{n-1})\delta_{n,m}, \qquad n, m < N. \tag{14}$$

However, in the continuous limit functions $\delta_n(x)$ and $\chi_n(x)$ should be replaced with distributions. One sees that $\mathrm{v}(x)$ demands using Dirac δ-function while $D(x)$ can be expressed through Heaviside θ-function. We find that

$$D(x) = p_1 + \sum_{n=1}^{N-1}\alpha_n\theta(|x| - a_n), \quad \mathrm{v}(x) = \mathrm{sign}(x)\sum_{n=1}^{N-1}\alpha_n\delta(|x| - a_n), \tag{15}$$

where $\mathrm{v}(x) = \partial_x D(x)$ and $\theta(x) = [1 + \mathrm{sign}(x)]/2$.

Therefore, (9) is reduced finally to

$$\partial_t P(x,t) = \partial_x[D(x)\partial_x P(x,t)], \qquad P(x,0) = \delta(x). \tag{16}$$

Although $D(x)$ contains information on geometry and phases of environment as "input", PDF is evolving in space-time and determines a normalized statistical measure μ_t for a fixed t:

$$\mathrm{d}\mu_t = P(x,t)\mathrm{d}x, \qquad \int \mathrm{d}\mu_t = 1. \tag{17}$$

In practice, one can often compute the probability of finding a walker at point x at time t,

$$\mathrm{Pr}(x,t) = \frac{1}{t}\int_0^t P(x,\tau)\mathrm{d}\tau, \tag{18}$$

that is the frequency of visiting a point x in a time t.

3. Finding a Solution

Developing an approach with $p_n > 0$, let us consider the functions

$$D(\alpha, x) = \sum_{n=1}^{N} (p_n)^\alpha \chi_n(x), \tag{19}$$

which are generated by the diffusivity function $D(x) \equiv D(1, x)$ and depend on both x and $\{p_n\}$ simultaneously, what requires specific rules of differential calculus.

Due to orthogonality of basis $\{\chi_n(x)\}$, one has

$$D(0, x) = 1, \qquad D(\alpha, x)D(\beta, x) = D(\alpha + \beta, x). \tag{20}$$

Differentiation (integration) of $D(\alpha, x)$ in space x is mainly related with a change of the basis, while exponents of parameters $\{p_n\}$ stay unchanged. One has

$$D'(\alpha + \beta, x) = D'(\alpha, x)D(\beta, x) + D(\alpha, x)D'(\beta, x), \tag{21}$$

where prime symbol means differentiation with respect to coordinate.

Numerical coefficients in the last formula are subjects of relation:

$$\begin{aligned}(p_{n+1})^{\alpha+\beta} - (p_n)^{\alpha+\beta} = {} & \frac{1}{2}\left[(p_{n+1})^\alpha - (p_n)^\alpha\right]\left[(p_{n+1})^\beta + (p_n)^\beta\right] \\ & + \frac{1}{2}\left[(p_{n+1})^\alpha + (p_n)^\alpha\right]\left[(p_{n+1})^\beta - (p_n)^\beta\right],\end{aligned} \tag{22}$$

where $1/2$ and sign "+" in brackets result from θ-function in D, while "−" results from D' containing δ-function.

To find PDF, we first concentrate the geometrical data in the new coordinate [13]

$$y(x) = \int_0^x D(-1/2, x')\mathrm{d}x'. \tag{23}$$

Although there exists one-to-one correspondence between x and y, the derivatives of $y(x)$ are singular, in general, at the points $x = \{\pm a_n\}$.

Integrating (23), we obtain that $y(x) = \mathrm{sign}(x)Y(-1/2, |x|)$, where

$$Y(\alpha, x) \equiv \frac{1}{2}\sum_{n=1}^{N} (p_n)^\alpha \left(a_n - a_{n-1} + |x - a_{n-1}| - |x - a_n|\right), \quad x \geq 0. \tag{24}$$

At this stage, let us define the new functions $\tilde{D}(\alpha, y(x)) = D(\alpha, x)$:

$$\tilde{D}(\alpha, y) = \sum_{n=1}^{N} (p_n)^\alpha \tilde{\chi}_n(y), \tag{25}$$

where $\tilde{\chi}_n(y) = \theta(|y| - b_{n-1}) - \theta(|y| - b_n)$ and $b_n \equiv y(a_n)$.

Then, introducing distribution $\mathcal{P}(y, t)$, we re-write statistical measure as

$$\begin{aligned}\mathrm{d}\mu_t &= \mathcal{P}(y, t)\mathrm{d}y \\ &= \mathcal{P}(y(x), t)\tilde{D}(-1/2, y(x))\mathrm{d}x.\end{aligned} \tag{26}$$

Substituting the re-defined $P(x, t)$ into (16), we have

$$\partial_t \mathcal{P} - \partial_y^2 \mathcal{P} = \kappa \partial_y(\beta \mathcal{P}), \qquad \mathcal{P}(y, 0) = \delta(y), \tag{27}$$

where constant κ regulates an interface effect between zones.

The right hand side of (27) can be reduced to the form with $\beta(y) = \tilde{D}(-1, y)\partial_y \tilde{D}(y)$. Computations lead to expressions:

$$\beta(y) = \mathrm{sign}(y) \sum_{n=1}^{N-1} \beta_n \delta(|y| - b_n), \qquad \beta_n = \frac{(p_{n+1})^2 - (p_n)^2}{2p_n p_{n+1}}, \tag{28}$$

$$B(y) = \sum_{n=1}^{N-1} \beta_n \theta(|y| - b_n), \qquad \partial_y B(y) = \beta(y). \tag{29}$$

Taking into account the specific properties of D-functions which should be investigated in more details, the form of β_n can vary what influences also the value of κ. However, a sign of β_n is defined by difference $p_{n+1} - p_n$. Moreover, $B(y)$ is not logarithmic function here, although $\beta(y)$ is similar to its log-derivative by construction.

We have already found that the bulk diffusion mainly depends on $y(x)$. Remaining problem is to account for the surface effect determined by $\beta(y)$ and κ. We attempt to find $\mathcal{P}(y, t)$ within the perturbative scheme, when

$$\mathcal{P}(y, t) = \varphi(y, t) + \sum_{r=1}^{\infty} \kappa^r \mathcal{S}_r(y, t), \tag{30}$$

where

$$\varphi(y, t) = \frac{1}{\sqrt{4\pi t}} \exp\left(-\frac{y^2}{4t}\right) \tag{31}$$

is Gaussian distribution resulting in probability (18):

$$\Phi(y, t) = \frac{1}{\sqrt{\pi t}} \exp\left(-\frac{y^2}{4t}\right) + \frac{|y|}{2t}\left[\mathrm{erf}\left(\frac{|y|}{2\sqrt{t}}\right) - 1\right]. \tag{32}$$

A possible solution of inhomogeneous equation $\partial_t \mathcal{S}_1 - \partial_y^2 \mathcal{S}_1 = \partial_y(\beta\varphi)$ is the function $\mathcal{S}_1(y, t) = -B(y)\varphi(y, t) - \mathcal{F}(y, t)$, where

$$\mathcal{F}(y, t) = -\frac{1}{2} \sum_{n=1}^{N-1} \beta_n [\varphi(|y - b_n| + b_n, t) + \varphi(|y + b_n| + b_n, t)]. \tag{33}$$

Finding $\mathcal{F}(y, t)$ from the auxiliary equation $\partial_t \mathcal{F} - \partial_y^2 \mathcal{F} = \beta \partial_y \varphi$, we contract $\beta\partial_y\varphi$ and φ playing a role of fundamental solution to diffusion equation with the use of formula:

$$\int_0^t \exp\left(-\frac{y_1^2}{4(t-\tau)} - \frac{y_2^2}{4\tau}\right) \frac{\mathrm{d}\tau}{\sqrt{(t-\tau)\tau^3}} = \sqrt{\frac{4\pi}{y_2^2 t}} \exp\left[-\frac{(|y_1| + |y_2|)^2}{4t}\right]. \tag{34}$$

Limiting ourselves by accounting for the first-order correction, we obtain our main result for PDF:

$$P(x, t) = \varphi(y(x), t)[1 - \kappa B(y(x))] D(-1/2, x) + \frac{\kappa}{2} \sum_{n=1}^{N-1} \beta_n [\varphi(|y(x) - b_n| + b_n, t) + \varphi(|y(x) + b_n| + b_n, t)] D(-1/2, x). \tag{35}$$



Figure 1. Probability to find a walker at point $x \in \mathbb{Z}$ for $t = 200$ (the total number of steps) in a triple-zone environment. Solid line is obtained with the use of (35), while the dashed line gives the result of averaging over 3000 random trajectories. Left panel: $2p_1 = 0.5$, $2p_2 = 0.9$, $2p_3 = 0.7$. Right panel: $2p_1 = 0.7$, $2p_2 = 0.5$, $2p_3 = 0.9$. In the both cases, $a_1 = 6$, $a_2 = 8$.

This formula looks applicable at $|\beta_n| < 1$ because the correction is of the same order of magnitude as φ. The PDF of ordinary RW is restored at $2p_n = 1$ for all n, when $\beta_n = 0$ and $y(x) = \sqrt{2}x$.

Of course, the number of independent zones decreases if the neighbouring zones have the same value of p_n and, therefore, are joint. Interface point $x = a_n$ between zones with p_n and $p_{n+1} = p_n$ becomes regular.

Note that (35) is similar to the amplitude squared in WKB approximation of quantum mechanics, and the points $x = \{\pm a_n\}$ are connected with the turns, induced by a potential in Schrödinger equation. In principal, a "potential" [13] in our model could appear by excluding term $\partial_y \mathcal{P}$ in (27) and would be determined by Schwarzian derivative of function (A.2)

$$f(y) = \int_0^y \tilde{D}(y')\mathrm{d}y'. \tag{36}$$

In order to get expression for probability $\mathrm{Pr}(x,t)$, we should simply replace function $\varphi(y,t)$ with $\Phi(y,t)$ in (35). To test our buildings, we compare the obtained $\mathrm{Pr}(x,t)$ with numerical simulations. Data are presented in figure 1. Analytical solution at $\kappa = 1$ is in agreement here with RW developing in triple-zone environment. However, one needs to account for highest-order corrections to reproduce better the considerable changes of the probability profile at short Δx.

4. Variance

To investigate the walker diffusion, let us calculate the variance $\langle x^2 \rangle$, that is,

$$\Lambda(t) \equiv \int x^2 \mathrm{d}\mu_t; \qquad \langle x \rangle \equiv \int x \mathrm{d}\mu_t = 0. \tag{37}$$

Note that $\Lambda(t)$ is linear in t in the case of purely Gaussian distribution.

In our case, one has $\Lambda(t) = \Lambda_0(t) + \kappa\Lambda_1(t) + O(\kappa)$, where

$$\Lambda_0(t) = \int_{-\infty}^{\infty} [x(y)]^2 \varphi(y,t)\mathrm{d}y, \quad \Lambda_1(t) = \int_{-\infty}^{\infty} [x(y)]^2 \mathcal{S}_1(y,t)\mathrm{d}y. \tag{38}$$

Although function $x(y)$ can be presented similarly to (24), it is usefully to re-write it in integrals in terms of $\{\tilde{\chi}_n(y)\}$. We find that

$$[x(y)]^2 = y^2 A_2(y) + |y|A_1(y) + A_0(y), \qquad A_s(y) = \sum_{n=1}^{N} A_{s,n}\tilde{\chi}_n(y); \tag{39}$$

where numeric coefficients are

$$A_{2,n} = p_n, \quad A_{1,n} = 2(\sqrt{p_n}c_n - p_n b_{n-1}), \quad A_{0,n} = \left(c_n - \sqrt{p_n}b_{n-1}\right)^2, \tag{40}$$

$$c_n = \sum_{m=1}^{n-1} \sqrt{p_m}(b_m - b_{m-1}). \tag{41}$$

Note that $A_2(y)$ coincides with diffusivity $\tilde{D}(y)$.

Combining the environment parameters $A_{s,n}$ and time dependent integrals, we write down the variance components:

$$\Lambda_0(t) = 2\sum_{s=0}^{2}\sum_{n=1}^{N} A_{s,n}[U_s(b_n,t) - U_s(b_{n-1},t)], \tag{42}$$

$$\begin{aligned}\Lambda_1(t) = & -2\sum_{s=0}^{2}\sum_{n=2}^{N} A_{s,n}B_n[U_s(b_n,t) - U_s(b_{n-1},t)] \\ & + \sum_{s=0}^{2}\sum_{n=1}^{N}\sum_{m=1}^{N-1} A_{s,n}\beta_m[V_s(b_n,t|b_m) - V_s(b_{n-1},t|b_m)],\end{aligned} \tag{43}$$

where $B_n = \sum_{m=1}^{n-1}\beta_n$ for $n \geq 2$, and

$$U_s(b,t) \equiv \int_0^b y^s \varphi(y,t)\mathrm{d}y, \tag{44}$$

$$V_s(B,t|b) \equiv \int_0^B y^s[\varphi(|y-b|+b,t) + \varphi(|y+b|+b,t)]\mathrm{d}y. \tag{45}$$

The Gaussian integrals yield

$$\begin{aligned}&U_0(b,t) = \frac{1}{2}\mathrm{erf}\left(\frac{b}{2\sqrt{t}}\right), \quad U_1(b,t) = \sqrt{\frac{t}{\pi}}\left[1 - \exp\left(-\frac{b^2}{4t}\right)\right], \\ &U_2(b,t) = t\ \mathrm{erf}\left(\frac{b}{2\sqrt{t}}\right) - b\sqrt{\frac{t}{\pi}}\exp\left(-\frac{b^2}{4t}\right).\end{aligned} \tag{46}$$

Using $U_s(b,t)$, we also get

$$\begin{aligned}V_0(B,t|b) = & U_0(B+2b,t) - \theta(b-B)U_0(2b-B,t) \\ & + \theta(B-b)\left[U_0(B,t) - 2U_0(b,t)\right], \\ V_1(B,t|b) = & U_1(B+2b,t) - 2U_1(2b,t) + 4bU_0(2b,t) - 2bU_0(B+2b,t) \\ & + \theta(b-B)\left[U_1(2b-B,t) - 2bU_0(2b-B,t) + 2bU_0(b,t)\right]\end{aligned} \tag{47}$$



Figure 2. Left panel: Time dependence of effective diffusivities $\Lambda/2t$ (solid line) and $\partial_t\Lambda/2$ (dashed line) for two sets of parameters as in figure 1. Right panel: Dependence of $C \equiv \partial_t^2\Lambda/2$ on t. Gray curves are for model with $p_1 = 0.35$, while black ones are for $p_1 = 0.25$.

$$+ \theta(B-b)U_1(B,t), \tag{48}$$

$$\begin{aligned} V_2(B,t|b) = {} & U_2(B+2b,t) - 2U_1(2b,t) - 4bU_1(B+2b,t) \\ & + 4b^2U_0(B+2b,t) \\ & - \theta(b-B)\left[U_2(2b-B,t) - 4bU_1(2b-B,t)\right. \\ & \left.+4b^2U_0(2b-B,t)\right] \\ & + \theta(B-b)\left[U_2(B,t) - 2U_2(b,t) + 4bU_1(b,t)\right. \\ & \left.-4b^2U_0(b,t)\right], \end{aligned} \tag{49}$$

where $\theta(0) = 1/2$ is assumed.

Analyzing $\Lambda(t)$, let us introduce effective diffusivities $\Lambda(t)/2t$ and $\partial_t\Lambda(t)/2$. Thus, if $p_n = p$ for all n, these coincide with p.

We also consider the second-order derivative of $\Lambda(t)$, denoted by C and connected with velocity autocorrelation function accordingly to Kubo's relation [14, 15]:

$$\frac{1}{2}\frac{\mathrm{d}\langle x^2(t)\rangle}{\mathrm{d}t} = \int_0^t \langle \dot{x}(t')\dot{x}(0)\rangle \mathrm{d}t', \qquad t \to \infty, \tag{50}$$

where $x(t)$ is the continuous stochastic coordinate with $\langle x(t)\rangle = 0$ and $\langle \dot{x}(t)\rangle = 0$.

The results based on our formulae are sketched in figure 2. We see on the left panel that there are the time intervals where the effective diffusivities are decreasing and increasing. These may correspond, respectively, to regimes of subdiffusion and superdiffusion, represented by $C < 0$ and $C > 0$ on the right panel. We conclude that a role of n-th zone in the diffusion picture depends mainly on adsorption effect which is proportional to probability $r_n = 1 - 2p_n$. Moreover, the model situations under consideration demonstrate inaccessibility of the regime with p_3 in a given time interval. Although, putting $b_N \to \infty$ before tending t to infinity, it is expected that the only terms with $U_2(b_N,t)/t \to 1$ should survive at $t \to \infty$ in a strong mathematical sense. However, we found that this limit is slowly reached because of environment structure.

In other words, there is always a non-vanishing probability of walker capture by n-th zone with $r_n > r_N$. This determines the time delay which may be estimated. Moreover, we assume an existence of different time scales which should be accurately defined [16] for a deeper understanding.

On the other hand, the right panel of figure 2 illustrates that the correlation C is relaxed with time, and information about early stage of evolution, developed in region near the origin, is lost. Being in infinitely wide N-th zone for a long time, a role of whole zone structure decreases in a walker history.

It is interesting to note that stage with $C < 0$ is often connected with a presence of liquid-like state in many-particle system [15]. This state emerges here due to zone with $2p = 0.5$ and could be intuitively foreseen, although sufficient conditions of its appearance are still unclear for us. We can only say that the zones with relatively small p_n keep a walker in a trap for a short time, so after a while the walker is able to diffuse away.

5. Discussion

Initially, we have formulated the model of one-dimensional RW with space dependent transition probability which is the same for the left and right steps almost everywhere. We describe a wandering in space with the finite-sized zones having the properties of adsorption in the bulk and reflection at the boundary points. It generalizes a problem of RW with various barriers [5, 6, 10] and allows us to speak about heterogeneous environment. Locating the N zones $x \in (-a_n, -a_{n-1}) \cup (a_{n-1}, a_n)$ densely along coordinate axis, transition probability in n-th zone is determined by constant parameter p_n, except the separation points where its value changes. This assumption is crucial for finding an analytical solution to the model equations in diffusion approximation at the large number of walker steps. Step-like dependence of diffusion coefficient on x prompts us to introduce the new coordinate $y(x)$, in terms of which the probability distribution is constructed with the use of Gaussian one (35). Nevertheless, the obtained distribution function is non-Gaussian because of complexity of environment and surface effects at singular points $\{\pm a_n\}$ where the derivatives of functions with respect to x diverge.

It seems to be difficult to describe analytically an effect of infinitely thin zone (a wall) without additional suggestions. Similar problem arises in zone interface description. Comparing the numerical RW simulations with discrete $x \in \mathbb{Z}$ and analytical solutions in continuous limit, we might effectively displace a position a_n of point-like barrier up to $a_n \pm 1/2$ in order to obtain the finitely wide border. Another possibility is to use the solutions with a gap [11] which are not considered here.

Actually, our model describes the diffusion process, taking into account the long-range properties of environment and neglecting local fluctuations $\delta D(x) \equiv D(x) - \bar{D}(x)$ of smooth-varied diffusivity $D(x)$, where

$$\bar{D}(x) = \sum_{n=1}^{N} D_n \chi_n(x), \qquad D_n = \frac{1}{2(a_n - a_{n-1})} \int_{-\infty}^{\infty} D(x)\chi_n(x)\mathrm{d}x. \tag{51}$$

Analyzing the results presented in figure 1, we can clarify the physical reasons of non-trivial distribution of the particles emanating from the origin.

Explanation of remarkable probability changes is based on a role of walker adhesion, which is forced by environment and related with probability $r_n = 1 - 2p_n$. Sequence of zones with different values of p_n is also significant: the sign of difference $p_{n+1} - p_n$ is responsible for preferable direction of the walk on interface of neighboring zones indexed by n and $n+1$. Relative time of the walker being in n-th zone depends on the number n, width $a_n - a_{n-1}$, and r_n. Thus, adhesion effect can be evaluated by characteristic time and length. Nevertheless, a long-time tail of the variance corresponds to the ordinary diffusion with $\langle x^2 \rangle \sim t$ at $t \to \infty$.

Alternative treatment can be done in terms of mechanics by introducing an effective walker "mass" $m_n = 1/2p_n$ for each zone of environment. Considering a macroscopic ensemble of the particles, we can interpret the mass changes as the result of geometrically dependent interaction among particles which is not specified in our picture but leads to appearance of the different states or phases confined within the given intervals.

It is expected that the Fourier transform $\tilde{P}(k,t)$ of PDF can give us the one-particle distribution function in momentum space k, accounting for the phase transitions in our case, in accordance with ideas of Bloch and Balescu. Indeed, replacing t and p_n with inverse temperature $1/T$ and $1/2m_n$, respectively, PDF of ordinary diffusion process,

$$\tilde{P}(k,t) \propto \mathrm{e}^{-k^2 pt}, \tag{52}$$

plays a role of Boltzmann distribution, which is a starting point of deriving thermodynamic functions and relations for many-particle ensemble.

Acknowledgments

A.N. thanks to A.M. Gavrilik (BITP) for fruitful discussions and to Yu.M. Sinyukov (BITP), suggesting us to highlight some aspects. A.N. is indebted for partial support of this work by Department of Physics and Astronomy of NAS of Ukraine.

Appendix A. Potential Model

Let us substitute $\mathcal{P}(y,t) = \exp\left[-B(y)/2\right]\Pi(y,t)$ in (27) at $\kappa = 1$. One gets

$$-\partial_t \Pi = -\partial_y^2 \Pi + V\Pi. \tag{A.1}$$

Potential V can be presented as

$$V(y) \equiv -\frac{1}{2}\left(\beta' - \frac{1}{2}(\beta)^2\right) = -\frac{1}{2}\mathrm{S}f(y), \tag{A.2}$$

where $\beta(y)$ and $f(y)$ are given by (28) and (36), respectively. We use formally notation for Schwarzian derivative $\mathrm{S}f \equiv (f''/f')' - (f''/f')^2/2$; and relation $\beta = f''/f'$.

Note that the form of (A.2) is similar to the potential of super-symmetric problem [17]. Function $u(y) \equiv 1/\sqrt{f'} = \tilde{D}(-1/2, y)$ is a solution to stationary equation

$$u'' + \frac{1}{2}(\mathrm{S}f)u = 0. \tag{A.3}$$

[1] See e.g. Shlesinger M F and West B (ed) *Random Walks and their Applications in the Physical and Biological Sciences* (AIP Conf Proc vol 109) (New York: AIP 1984); Spitzer F *Principles of Random Walk* (Berlin: Springer 1976)
[2] Havlin S and Abraham D Ben 1987 *Phys. Adv.* **36** 695
[3] Liu L, Li P and Asher S A 1999 *Nature* **397** 141
[4] Lehner G 1963 *Ann. Math. Statist.* **34** 405
[5] Gupta H S 1966 *J. Math. Sci.* **1** 18
[6] Percus O E and Percus J K 1981 SIAM *J. Appl. Math.* **40** 485; Percus O E 1985 *Adv. Appl. Prob.* **17** 594
[7] Tanner J E 1978 *J. Chem. Phys.* **69** 1748
[8] Burada P S, Hänggi P, Marchesoni F, Schmid G and Talkner P 2009 *Chem. Phys. Chem.* **10** 45
[9] Novikov D S, Fieremans E, Jensen J H and Helpern J A 2011 *Nat. Phys.* **7** 508
[10] Ascher M 1960 *Math. Comp.* **14** 346
[11] Powels J G, Mallett M J D, Rickayzen G and Evans W A B 1992 *Proc. R. Soc. Lond.* A **436** 391
[12] Vanderzande C *Lattice models of polymers* (Cambridge University Press 1998)
[13] Güngör F 2015 Equivalence and Symmetries for Linear Parabolic Equations and Applications Revisited arXiv:1501.01481 [math-ph]
[14] Kubo R 1966 *Rep. Prog. Phys.* **29** 255
[15] Brodin A, Turiv T and Nazarenko V 2014 *Ukr. J. Phys.* **59** 775
[16] Dechant A, Lutz E, Kessler D A and Barkai E 2014 *Phys. Rev.* X **4** 011022
[17] Bravo R and Plyushchay M S 2016 *Phys. Rev.* D **93** 105023